


CARESS Working Paper #95-15
An "Anti-Folk Theorem" for a Class of Asynchronously
Repeated Games

by

Roger Lagunoff and Akihiko Matsui



UNIVERSITY of PENNSYLVANIA

Center for Analytic Research
in Economics and the Social Sciences

McNEIL BUILDING, 3718 LOCUST WALK
PHILADELPHIA, PA 19104-6297

CARESS Working Paper #95-15
An “Anti-Folk Theorem” for a Class of Asynchronously
Repeated Games

Roger Lagunoff* Akihiko Matsui†

August 12, 1995‡

Abstract

It is well known from the Folk Theorem that infinitely repeated games admit a multitude of equilibria. This paper demonstrates that in some types of games, the Folk Theorem form of multiplicity is an artifact of the standard representation which assumes perfect synchronization in the timing of actions between the players. We define here a more general family of repeated settings called *renewal games*. Specifically, a renewal game is a setting in which a stage game is repeated in continuous time, and at certain stochastic points in time determined by an arbitrary renewal process some set of players may be called upon to make a move. A stationary, ergodic Markov process determines who moves at each decision node. We restrict attention in this paper to a natural subclass of renewal games called *asynchronously repeated games*, in which no two individuals can change their actions simultaneously. Special cases include the alternating move game, and the Poisson revision game. In the latter, each player adjusts his action independently at Poisson distributed times.

Our main result concerns asynchronously repeated games of pure coordination (where the payoffs of all players in the stage game are identical up to an affine transformation): given $\epsilon > 0$, if players are sufficiently patient then every Perfect equilibrium payoff comes within ϵ of the Pareto dominant payoff. We also show that the “Folk wisdom” in the standard model that repetition always expands (weakly) the set of equilibrium payoffs is not true generally in asynchronously repeated games.

Journal of Economic Literature Classification Numbers: C72, C73.

Keywords: repeated games, asynchronously repeated games, renewal games, coordination games.

Department of Economics, University of Pennsylvania, Philadelphia, PA 19104, and Department of Economics, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061.

Department of Economics, University of Pennsylvania, Philadelphia, PA 19104, and Department of Economics, University of Tsukuba, Ibaraki 305, Japan.

We would like to thank Eddie Dekel, Hseuh-ling Huyn, Mamoru Kaneko, Peter Linhart, George Mailath, Stephen Morris, Roy Radner, Phil Reny, Masahiro Okuno-Fujiwara, and numerous seminar participants for many useful discussions and suggestions.

1. Introduction

The theory of infinitely repeated games offers little predictive content. It is well known from the Folk Theorem that infinitely repeated games admit a multitude of equilibria.¹ This paper demonstrates that in some types of games, the form of multiplicity given in the Folk Theorem is intimately connected with the way that repeated play is modelled in the standard repeated game formulation.

To focus on a particularly robust multiplicity problem, it is known that each one shot Nash payoff is an equilibrium payoff of the infinite repetition of a stage game, and each payoff derived as a convex combination of one-shot equilibrium payoffs (both pure and mixed equilibria) is also an equilibrium payoff of the repeated game. In the simple, canonical pure coordination game G_1 below, this means that any symmetric payoff pair (u, u) with u in the interval $[2/3, 2]$ is an equilibrium payoff of the infinite repetition of G_1 . These payoffs are robust with respect to variations in both discount parameters and the equilibrium concept. That is, these payoffs remain equilibrium payoffs for all discount parameters, and they remain perfect equilibrium payoffs as well.

$$\begin{array}{c}
 \mathbf{2} \\
 \begin{array}{cc}
 & \begin{array}{cc} s_2^* & \bar{s}_2 \end{array} \\
 \begin{array}{c} \mathbf{1} \\ s_1^* \\ \bar{s}_1 \end{array} & \begin{array}{|cc|} \hline 2, 2 & 0, 0 \\ \hline 0, 0 & 1, 1 \\ \hline \end{array} \\
 & \mathbf{G}_1
 \end{array}
 \end{array}$$

These observations are neither new nor surprising. What we emphasize here is the fact that repetition always *expands* (weakly) the set of equilibrium outcomes, and that this is a particularly robust phenomena — *within the standard formulation of repeated play*. Our contribution here is to demonstrate why this last addendum is necessary.

The standard model of repeated strategic play is a discretely repeated, simultaneous move game. This formulation assumes a perfect synchronization in the timing of actions between the players. Alternatively, this assumption may be interpreted as having each player move in ignorance of the other players' current move. The effect of this assumption in game G_1 above, for example, is that each player may take an action consistent with a Pareto inferior static equilibrium only because he expects that all others will do the same. Since all players move at once, no player can unilaterally signal his intent to do otherwise. While the synchronized moves is not an unreasonable model of repetition in certain settings, it is not clear why it should *necessarily* be the benchmark setting for repeated play, especially in light of the multiplicity dilemma. It seems natural in many contexts that players move asynchronously to initiate a unilateral move. While signalling can occur in the standard model, it remains part of the coordination built in to equilibrium beliefs rather than a

¹See, for example, Abreu (1988) or Fudenberg and Maskin (1986). References to earlier results may be found in a survey by Aumann (1981).

physical attribute of the game. If no two players can move at once then even if a player expects that other players will choose the inferior equilibrium he must wait for them to actually do so since actions cannot be perfectly synchronized. The lack of synchronization may therefore eliminate a “coordinated mistake.”

In this paper we describe a general class of games with repeated interaction, some of which break the perfect synchronization of the standard model. In so doing we are able to significantly pare down the set of equilibria in certain types of such games. We call games in this class *renewal games*. Specifically, a renewal game is defined as a setting in which a stage game is repeated in continuous time, and at certain stochastic points in time, determined by an arbitrary renewal process, some set of players may be called upon to make a move. A stationary Markov process operates on a set of “decision states,” each decision state determining “who moves” at each jump in the renewal process. The process is assumed to have the property that every player will obtain a chance (almost surely) to make a move after any state. The standard repeated game model is shown to be a special case of a renewal game.

Though the characterization of equilibrium payoffs for general renewal games is our eventual goal, we will limit our attention in this paper to an interesting subclass of renewal games in which no more than one player may revise his action at any given time. We call the games that fit this description *asynchronously repeated games*.

Two special cases of asynchronously repeated games are the two player alternating move game (see, for example Maskin and Tirole (1988)), and the Poisson revision game studied in several evolutive models (see Lagunoff and Matsui (1995) for one formal description). In the former, the renewal process is deterministic, and each player moves at every other decision state without moving together with the other player. In the latter, each player can adjust his action at stochastic intervals determined by a Poisson process. Each player’s adjustment process is independent of any other player’s. Both examples share the general characteristic of asynchronously repeated games which is that no two individuals can change their actions simultaneously. In such games the players’ adjustments may exhibit some inertia due to each player’s (possibly stochastic) delay. We consider equilibria which are robust to taking the limit as the players’ discount rates approach zero, or alternatively, the average delay between a player’s revision opportunities goes to zero. In all such games, we do not bound the complexity of each player’s behavior strategy except to assume that strategies are conditioned on the history of behavior at decision opportunities.

Our main result pertains to pure coordination stage games — games in which payoffs of all players are identical (up to an affine transformation). We show that if players are sufficiently patient or, equivalently, if players can revise sufficiently quickly, then for any $\epsilon > 0$ every Perfect equilibrium payoff comes within ϵ of the Pareto dominant payoff. This result starkly contrasts with the Folk Theorem, as it essentially rules out all the inefficient payoffs in the above example, particularly the inefficient static Nash equilibrium (hence the term, “anti-Folk Theorem.”) This result suggests that teams will solve their coordination problems in asynchronously repeated games.

Additionally, for a fixed discount rate close to zero we can find a neighborhood in payoff space of the pure coordination game in which the “anti-Folk Theorem” holds. Unfortunately, the uniqueness result seems to be limited to stage games close (in the sense just described) to the pure coordination game. We give an example in which the multiplicity of PE persists for patient players if costs of miscoordination are highly asymmetric. However, we show that there is a neighborhood of discount rates that support the Pareto dominant outcome as the unique PE payoff in a neighborhood of asynchronously repeated games. Again, this contrast with the standard model since inefficient Nash payoffs of the stage game are not sustained in these neighborhoods of repeated games and discount rates.

Finally, we establish a “continuity result” for a one parameter family of renewal games. The standard repeated model is determined by one extremal value of the parameter, while an asynchronously repeated game is determined by the other. At issue is whether the “Anti-Folk Theorem” represents a discontinuity when the standard model is perturbed. We show that the set of PE payoffs shrinks continuously from the set of Folk Theorem payoffs to the Pareto dominant singleton as the parameter varies between the standard and the asynchronous models, resp.

We emphasize that models of asynchronous interaction are not new. In differing contexts, Gale (1995), Morris (1995), Perry and Reny (1993), Maskin and Tirole (1988), Farrell and Saloner (1985), and our previous work (some with Rafael Rob),² as well as others have studied consequences of asynchronous choice. In Section 6 we relate these to the present paper, and we look at other literature which contrasts with the Folk Theorem for repeated games. To be clear, the present analysis is intended to be more suggestive than definitive. While a complete characterization is preferable, our purpose here is to give some support for the study of a more general class of repeated interactions and to suggest that such an analysis will yield, at times strikingly, different results.

The paper proceeds as follows. Section 2 describes the model and defines the asynchronously repeated game and the equilibrium concept of Markov perfection. Section 3 states and proves the uniqueness results for games of pure coordination. Section 4 discusses games of impure coordination. We show how the results may be sensitive to asymmetries in the costs of miscoordination. We provide a genericity result which describes the sense in which the results are robust to perturbations in the pure coordination structure. We also show that there are discount factors under which the “anti-Folk Theorem” holds for a more general class of coordination games. Section 5 gives the continuity result, and Section 6 discusses related literature.

2. A Model of Asynchronously Repeated Interaction

2.1. Stage Game

Let $G = (I, (S_i)_{i \in I}, (u_i)_{i \in I})$ denote a normal form stage game where I is the finite set of players,

²These include Matsui and Rob (1992), Lagunoff and Matsui (1995), and Lagunoff (1995).

S_i ($i \in I$) is the finite set of actions for player i , and $u_i : S \equiv \times_{i \in I} S_i \rightarrow \mathfrak{R}$ is the payoff function for player $i \in I$. Without loss of generality, assume that $S_i \cap S_j = \emptyset$ for all $i \neq j$. We will call an element of $s = (s_1, \dots, s_{|I|}) \in S$ a *behavior profile* (or simply “profile”). Given some $\hat{s}_i \in S_i$, let $s \setminus \hat{s}_i = (s_1, \dots, s_{i-1}, \hat{s}_i, s_{i+1}, \dots, s_{|I|})$. The tuple of payoff functions is denoted by $u = (u_i)_{i \in I}$. A mixed strategy for i will be denoted by σ_i and has the standard properties: $\sigma_i : S_i \rightarrow [0, 1]$ and $\sum_{s_i} \sigma_i(s_i) = 1$. A *mixed profile* is given by $\sigma = (\sigma_i)_{i \in I}$. Finally, a game G is a *coordination game* if its Nash equilibria are Pareto ranked and there is some Nash equilibrium that strictly Pareto dominates every other profile of the game.

2.2. Renewal Games and Asynchronously Repeated Games

In this Section, we introduce a framework that encompasses a wide variety of repeated strategic environments. Consider a continuous repetition of a stage game G . After the first choice at time zero, which is either chosen by the players as in standard repeated games, or chosen by nature as in models of evolution, all players’ decision points are determined by a renewal process and a Markov process with finitely many states. It will be clear that the following description and results will not be altered by specification of choice at time zero. In the following, decision nodes refer to the decision nodes other than that in the first stage (at time zero).

Formally, let X_1, X_2, \dots be an infinite sequence of i.i.d. nonnegative random variables which follow a (marginal) probability measure ν with $E_\nu(X_1) < \infty$ and $\nu(X_1 > \epsilon) > \delta$ for some $\epsilon, \delta > 0$. It is also assumed that $\nu(X_1 = 0) = 0$ so that the orderliness condition for the renewals is guaranteed. Then let $T_0 = 0$ and $T_k = T_{k-1} + X_k = X_1 + \dots + X_k$ ($k = 1, 2, \dots$). T_k is the time elapsed before the k th revision point. At each decision point a state ω is determined from a finite set Ω according to a Markov process $\{Y_k\}_{k=1}^\infty \in \Omega^\infty$ where $Y_k = \omega$ ($\omega \in \Omega$) implies that state ω is reached at time T_k . We denote $p_{\omega\omega'} = Pr(Y_{k+1} = \omega' \mid Y_k = \omega)$ for $\omega, \omega' \in \Omega$. Let $\Omega_i \subseteq \Omega$ denote the nonempty set of states in which player i has a decision node. Let $\Omega_0 \subseteq \Omega$ be the set of “inertial” states in which no player has a decision node. By definition, $\Omega_0 = \Omega - (\cup_{i \in I} \Omega_i)$. Note that $\Omega_i \cap \Omega_j$ ($i \neq j$) may or may not be empty. To summarize, the renewal process, ν , determines when the decision nodes (the “jumps”) occur, while the Markov transition, p , determines who moves at each node.

Definition 1. A *renewal game* is a tuple

$$\Gamma = \langle G, \nu, (p_{\omega\omega'})_{\omega, \omega' \in \Omega}, r \rangle,$$

where $r > 0$ is a common discount rate, and for all $\omega \in \Omega$ and all $i \in I$, there exists a chain of states, $\omega^0, \omega^1, \omega^2, \dots, \omega^M$ ($M < |\Omega|$), with $\omega = \omega^0$ and $\omega^M \in \Omega_i$ such that $p_{\omega^{m-1}\omega^m} > 0$ ($m = 1, \dots, M$) (from any state, every player obtains a revision node).

Since the number of states is finite, renewal games have the property that from any state, every player obtains revision nodes infinitely many times, and the expected time interval between two revision nodes is finite. Standard discounted repeated games are renewal games which may be described in several ways. One straightforward way is: $\Omega = \Omega_i, \forall i \in I$, and $\nu(X_1 = 1) = 1$. However, we wish to specialize further to only those renewal games in which choice is asynchronous.

Definition 2. An *asynchronously repeated game* is a renewal game in which $\Omega_i \cap \Omega_j = \emptyset$, $\forall i \neq j$

In asynchronously repeated games, no two individuals have simultaneous revision opportunities. When $\omega \in \Omega_i$, we will write $i(\omega) = i$. Some examples are:

Example 1 (alternating move game). Let $I = \{1, 2\}$, $\Omega = I$, $X_1 \equiv 1$, and $p_{ij} = 1$ if $i \neq j$. Then the decision points are deterministic, and players' revision nodes alternate.

Example 2 (Poisson revision process). Let $\Omega = I$. And let X_1 follow an exponential distribution with parameter $\lambda > 0$, i.e., $\nu(X_1 < x) = 1 - e^{-\lambda x}$, and let $p_{ij} = p_j$ for all $i, j \in I$. Then players' revision points are independent of each other, and player i 's decision points ($i \in I$) follow a Poisson process almost surely with parameter λp_i .

Example 3 (ϵ -approximation of the standard repeated game). Let $I = \{1, 2\}$, and

$$\Omega = \overbrace{\{\omega_0^1, \dots, \omega_0^M\}}^{\Omega_0} \cup \overbrace{\{\omega_1^1, \omega_1^2\}}^{\Omega_1} \cup \overbrace{\{\omega_2^1, \omega_2^2\}}^{\Omega_2}$$

Then assume that Y_k 's follow the process illustrated in Figure 2 below. In the figure, the process proceeds through the inertial states until ω_0^M . At that time the process moves to either player 1's or player 2's decision node with probability 1/2 each. Let $X_1 \equiv 1/(M+2)$. Then if M is sufficiently large, the process approximates the standard repeated game in the sense that each player has a revision opportunity once in a unit of time, and that the two players' decision nodes are very close in timing.




Figure 2: An ϵ -approximation of the standard repeated game.

2.3. Recursive Structure

One additional assumption that we will make will be to restrict in a natural way the class of behavior strategies that individuals use. We assume that individuals only condition on the *sequences* of decision points rather than on the time interval between them. This restriction significantly reduces the notational burden, and it turns out that it in no way alters main results of this paper.³

Restricting conditioning events to the “jumps” rather than on time intervals between jumps allows a straightforward recursive representation of individuals’ continuation values in the asynchronous model. To formulate this, let $s(t) = (s_1(t), \dots, s_{|I|}(t))$ denote the behavior profile at time t , and let $N_t \equiv \inf\{k : T_k > t\}$ denote the number of renewals up to time t . Due to the assumption on X_k ’s, $N_t < \infty$ holds almost surely. Then define the space of histories H such that a history $h^t \in H$ is given by $h^t = (y^t, s^t)$ wherever N_t is finite, and

$$y^t = \{Y_k\}_{k \leq N_t},$$

³The interested reader can obtain a note by Lagunoff and Matsui (1995) which proves Theorems 1 and 2 for the case of full-fledged behavior strategies.

and

$$s^t = \{s(T_k)\}_{k \leq N_t}.$$

The null history is denoted by h^0 . Since at most one player switches his action at a time after the initial profile we write $s^t = (s(h^0), s_{Y_1}, \dots, s_{Y_{N_t}})$ whenever convenient. We let $s(h)$ and $\omega(h)$ denote the current behavior and state at history h , resp., and let $i(h)$ denote the last player whose decision node was reached. We also define $h^{t-} = (y^t, \{s(T_k)\}_{k < N_t})$ so that h^{t-} includes the same information as history h^t except for the behavior profile at time t which may include a new decision by a player. We write H^- for the set of all such *conditioning histories* and denote an element by h^- .

A *strategy* for player i is a history contingent action given by the function $f_i : H^- \rightarrow \Delta(S_i)$. Although this formulation appears to require that i is unable to condition on the current behavior profile, recall that decisions are only made at the “jumps” in the renewal process.

The play of the game proceeds as follows. At time zero, all the players in I simultaneously take actions, $f_i(h^{0-})$ for $i \in I$. Suppose $T_1 = t_1 = x_1$ and $Y_1 = y_1 \in \Omega_i$. If $i = 0$, nothing changes except the Markov “clock.” If $i \in I$, then $h^{t_1-} = (f(h^{0-}); y_1)$, player i 's revision node is reached at time t_1 , and he takes action $f_i(h^{t_1-})$. History at time t_1 becomes $h^{t_1} = (y_1; f(h^{0-}), f(h^{0-}) \setminus f_i(h^{t_1-}))$. Given a strategy profile f and $h \in H^t$ (resp. $h^- \in H^{t-}$), $\{\tilde{s}(f|h)(\tau)\}_{\tau > t}$ (resp. $\{\tilde{s}(f|h^-)(\tau)\}_{\tau \geq t}$) denotes an induced (stochastic) path of action profile after h (resp. h^-).

Given a history $h^t \in H$ and a strategy profile $f = (f_i)_{i \in I}$, we define the conditional discounted expected payoff to player i at time t by

$$V_i(f|h^t) = \tau \int_t^\infty e^{-r(\tau-t)} E \left[u_i \left(\tilde{s}(f|h^t)(\tau) \right) \right] d\tau,$$

where $E[\cdot]$ is the expectation operator. A strategy profile f^* is called a *perfect equilibrium (PE)* if for each $i \in I$, f_i^* is a best response to $(f_j^*)_{j \neq i}$ after every history h^t , i.e.,

$$V_i(f^*|h^t) \geq V_i(f^* \setminus f_i|h^t)$$

for any of player i 's strategies f_i .

One immediate result in the asynchronously repeated game is that for almost all histories, mixed strategies will not be used at any revision opportunity. Hence, the mixed strategy minimax payoff which is always an equilibrium of the stage coordination game is not the benchmark here necessarily. Hereafter, we will often denote a pure strategy by $f_i(h^-)$ as well as a mixed strategy.

Given a history $h = h^t \in H$, let $h^t \circ (\omega; s_j)$ denote the concatenated history in which, after h^t , the next state $\omega \in \Omega_j$ is reached, at which player j takes s_j . Given $h^t \in H$, $h^t \circ \omega \in H^-$ is a path such that after h^t , state ω is reached (without specifying the revised action). Using this expression, the value after $h \circ \omega \in H^-$ is given by

$$V_i(f|h \circ \omega) \equiv V_i(f|h \circ (\omega; f_{i(\omega)}(h \circ \omega))).$$

The analysis will make extensive use of the following recursive formulation. The continuation value to i induced by f after history $h^t \in H$ with $\omega(h^t) = \omega$ may be expressed as

$$V_i(f|h^t) = (1 - \sum_{\omega' \in \Omega} \theta_{\omega\omega'})u_i(s(h^t)) + \sum_{\omega' \in \Omega} \theta_{\omega\omega'}V_i(f|h^t \circ \omega'). \quad (2.1)$$

where

$$\begin{aligned} \theta_{\omega\omega'} &\equiv p_{\omega\omega'} \int_0^\infty e^{-rt} d\nu(t) \\ &= \text{expected discounted probability that } \omega' \\ &\quad \text{is the first state reached from } \omega. \end{aligned} \quad (2.2)$$

2.4. Existence

The following is a standard proof for the existence of perfect equilibria. What is proven here is actually the existence of so called Markov perfect equilibrium.

Theorem 0 *For any asynchronously repeated game $\Gamma = \langle G, \nu, p, r \rangle$ there exists at least one perfect equilibrium.*

Proof Partition H^- into $\wp = \{H_{\omega s}^-\}_{s \in S, \omega \in \Omega}$ such that

$$H_{\omega s}^- = \{h^- \in H^- | \omega(h^-) = \omega, s(h^-) = s\}, \forall \omega \in \Omega, \forall s \in S.$$

Observe that \wp constitutes the ‘‘payoff relevant’’ set of states. Suppose that each player takes a $\sigma(\wp)$ -measurable behavior strategy where $\sigma(\wp)$ is the σ -algebra generated by \wp . Then the play of the game follows a Markov process, and we can represent a strategy of player i by the ‘‘Markovian’’ function $\psi_i \in [\Delta(S_i)]^\wp$. The strategy represented by ψ_i is denoted by f_{ψ_i} .

For each $i \in I$ let $BR^i = \{BR_{\omega s}^i\}_{\omega \in \Omega, s \in S}$ satisfy

$$BR_{\omega(h)s}^i(\psi) = \arg \max_{f_{\psi_i}} V(f_\psi \setminus f_{\psi_i} | h) \quad (2.3)$$

Letting $BR = (BR_i)_{i \in I}$, equation (2.3) defines an upper hemicontinuous correspondence $BR : \times_i [\Delta(S_i)]^\wp \rightarrow \times_i [\Delta(S_i)]^\wp$ where $\times_i [\Delta(S_i)]^\wp$ is compact and convex. Therefore, by Kakutani’s fixed point theorem, there exists ψ such that $\psi \in BR(\psi)$ holds. Standard arguments show that the corresponding strategy f_{ψ_i} is a best response to $(f_{\psi_j})_{j \neq i}$ within the class of all strategies after any history h (or, more precisely, after any h^-) since all j , ($j \neq i$) only vary their behavior over ‘‘states’’ $\omega s \in H_{\omega s}^-$. Hence, f_ψ is a perfect equilibrium. \square

3. Pure Coordination

A game G is a pure coordination game if $u_i = u_j = u$ for all i and j .⁴ Let s^* denote the profile that gives each player his highest payoff u^* . The first part of our main result states that, independently of the discount rate r , the unique continuation value after any history in which the current profile is s^* must give the optimal payoff $u_i^* = u_i(s^*)$ for all $i \in I$.

Theorem 1. *Given any asynchronously repeated game $\Gamma = \langle G, \nu, (p_{\omega\omega'}), r \rangle$ in which G is a pure coordination game, for any perfect equilibrium f of Γ , for any history $h \in H$ with $s(h) = s^*$, and for all $i \in I$, $V_i(f|h) = u_i^*$ holds.*

proof of Theorem 1. Observe first that due to the identical payoffs, we can drop the subscript i on u_i and continuation values V_i . Fix a perfect equilibrium f . Recall that $s(h)$ is the current action profile given history $h \in H^t$. Define $\underline{V} = \inf_{\{h: s(h)=s^*\}} V(f|h)$. This is the infimum value of the game when the current behavior profile is s^* . Since the payoff space is bounded from below, this infimum exists. Fix $\epsilon > 0$. Then there exists $h = h_\epsilon \in H^t$ for some $t > 0$ such that $s(h) = s^*$ and

$$\underline{V} > V(f|h) - \epsilon. \quad (3.1)$$

Recall from (2.1) the continuation value after h which is given by

$$V(f|h) = (1 - \sum_{\omega \in \Omega} \theta_{\omega(h)\omega}) u^* + \sum_{\omega \in \Omega} \theta_{\omega(h)\omega} V(f|h \circ \omega). \quad (3.2)$$

If $\omega \in \Omega_0$, then $s(h \circ \omega) = s^*$, and therefore, $V(f|h \circ \omega) \geq \underline{V}$. Observe also that since f is a perfect equilibrium strategy profile, it must be the case that for all $i \in I$ and all $\omega \in \Omega_i$,

$$V(f|h \circ \omega) \geq V(f|h \circ (\omega; s_i^*)) \geq \underline{V}$$

where the second inequality holds due to the definition of \underline{V} . Substituting these inequalities into (3.2) and using (3.1), we obtain

$$\underline{V} > (1 - \sum_{\omega' \in \Omega} \theta_{\omega(h)\omega'}) u^* + \sum_{\omega' \in \Omega} \theta_{\omega(h)\omega'} \underline{V} - \epsilon \quad (3.3)$$

Since $\sum_{\omega \in \Omega} \theta_{\omega(h)\omega} = \int_0^\infty e^{-r\tau} d\nu(\tau)$, (3.3) implies

$$\underline{V} > u^* - \frac{\epsilon}{1 - \int_0^\infty e^{-r\tau} d\nu(\tau)}.$$

Since ϵ is arbitrary and independent of r , $\underline{V} \geq u^*$ holds. \square

⁴This can be weakened so that we require only that $u_i = \alpha u_j + \beta$ for $\alpha > 0$ and $\beta \in \mathfrak{R}$.

A Remark on the Logic of Theorem 1. Before proceeding to the second part of the result (Theorem 2), we emphasize that since Theorem 1 (and Theorem 2 which follows) holds for any asynchronously repeated game of pure coordination, what matters for such games is the fact that choice is asynchronous rather than the specific form of asynchronous choice. Given the pure coordination structure, the logic utilizes standard arguments from dynamic programming. Figure 3 below is a “false counterexample” which further illustrates why asynchronous choice is so crucial. For exposition, suppose that Γ denotes an alternating move, asynchronously repeated version of the pure coordination game G_1 .




Figure 3: Equilibrium paths in an alternating move game.

In Figure 3 the first layer describes the candidate “equilibrium” path. Subsequent layers are “equilibrium continuations” after deviations. In “equilibrium” the players remain in s^* for a fixed time before player 1 chooses \bar{s}_1 , thus signalling a switch to the inferior equilibrium \bar{s} which lasts for

L_1 periods. Initially player 1, who is prescribed to be the first to switch to \bar{s}_i , may choose to deviate and remain in s^* . The second layer is a “punishment path” which prescribes a longer time length of L_2 periods spent in \bar{s} . Each successive layer lengthens the time in \bar{s} . The infimum \underline{V} defined in the proof is approximated by the average of the payoffs in the k th layer for k large. We argue that there is some k large enough such that a deviation — remaining in s^* — is profitable. The reason is simple. The average payoff in layer k converges to 1 as $L_k \rightarrow \infty$ as $k \rightarrow \infty$. Therefore, the punishment for remaining in s^* in layer k is the additional $L_{k+1} - L_k$ periods which lowers the average payoff by $(n(L_{k+1} - L_k))/(n + L_{k+1})(n + L_k)$ where n is the number of periods spent in s^* . Clearly this cost approaches zero in k . By contrast the benefit to deviating is 2Δ for $\Delta > 0$ which is independent of k . Once it is established that a player will deviate in the k th layer for k large enough, backward induction then destroys the equilibrium.

The second part of the main result is that any continuation value in a perfect equilibrium is arbitrarily close to the Pareto efficient value for sufficiently patient players.

Theorem 2. *Given any asynchronously repeated game $\Gamma = \langle G, \nu, (p_{\omega\omega'}), r \rangle$ in which G is a pure coordination game, and given any $\epsilon > 0$, there exists $\bar{r} > 0$ such that if $r \in (0, \bar{r})$, then for all perfect equilibrium f of Γ , for all histories $h \in H$, and for all $i \in I$, $V_i(f|h) > u_i^* - \epsilon$ holds.*

proof of Theorem 2. The proof will use a backward induction argument to establish that after any history the equilibrium continuation must be within ϵ of the maximal payoff u^* . As in the first theorem, we drop the subscript i . Fix $\epsilon > 0$. For each $k = 0, 1, \dots, |I|$, define

$$S^k = \{s \in S : |\{i \in I | s_i \neq s_i^*\}| = k\}.$$

S^k is the set of behavior profiles of the stage game that require k individuals to change their actions to get to profile s^* . By this definition, $S^0 = \{s^*\}$.

Fix a perfect equilibrium f . For each $s \in S$ and each $\omega \in \Omega$, we let $\underline{V}_\omega^s = \inf_{s(h)=s, \omega(h)=\omega} V(f|h)$, and $\underline{V}^s = \inf_{s(h)=s} V(f|h)$. Note that $\underline{V}^s = \min_{\omega \in \Omega} \underline{V}_\omega^s$.

The proof will proceed as follows: recall from Theorem 1 that after a history h with $s(h) \in S^0$, i.e., $s(h) = s^*$, the continuation value $V(f|h)$ is u^* . Given ϵ , the first step is to find a bound \bar{r} such that if $r < \bar{r}$ then for all $k = 1, 2, \dots, |I|$, there is some profile $s' \in S^{k-1}$ such that after any h with $s(h) \in S^k$, the continuation value satisfies

$$V(f|h) > \underline{V}^{s'} - \frac{\epsilon}{|I|}. \quad (3.4)$$

This turns out to be the crucial step. The choice of \bar{r} must be determined independently of f . The next step is to verify from backward induction on S^k that (3.4) holds for all such $s' \in S^{k-1}$. Finally, we verify that $0 < r < \bar{r}$ implies that for all history $h \in H$, $V(f|h) > u^* - \epsilon$. Thus, the proof will be completed.

First, we determine the value of \bar{r} independently of f . Recalling that the definition of $\theta_{\omega\omega'}$ is a

function of r , define the bound \bar{r} so as to satisfy

$$\min_{\substack{(\omega_0, \omega_1, \dots, \omega_N) \in \Omega^{N+1} \\ 2 \leq N \leq |\Omega|}} \left\{ \frac{\theta_{\omega_0 \omega_1} \cdots \theta_{\omega_{N-1} \omega_N}}{\theta(\omega_0, \dots, \omega_N)} : \theta_{\omega_0 \omega_1} \cdots \theta_{\omega_{N-1} \omega_N} > 0 \right\} > 1 - \frac{\epsilon}{2|I|} \frac{1}{u^* - \min_s u(s)}, \quad (3.5)$$

where

$$\begin{aligned} \theta(\omega_0, \dots, \omega_N) &= 1 - \sum_{\ell=1}^N \prod_{n=1}^{\ell-1} \theta_{\omega_{n-1} \omega_n} \sum_{\omega' \neq \omega_\ell} \theta_{\omega_\ell - 1 \omega'} \\ &= 1 - \sum_{\omega' \neq \omega_1} \theta_{\omega_0 \omega'} - \theta_{\omega_0 \omega_1} \sum_{\omega' \neq \omega_2} \theta_{\omega_1 \omega'} - \cdots - \theta_{\omega_0 \omega_1} \cdots \theta_{\omega_{N-2} \omega_{N-1}} \sum_{\omega' \neq \omega_N} \theta_{\omega_{N-1} \omega'}. \end{aligned}$$

Such an \bar{r} can be found since both $\theta_{\omega_0 \omega_1} \cdots \theta_{\omega_{N-1} \omega_N}$ and $\theta(\cdot)$ converge to $p_{\omega_0 \omega_1} \cdots p_{\omega_{N-1} \omega_N} > 0$ as r goes to zero. Also, let $\underline{\theta} = \min\{\theta(\omega_0, \dots, \omega_N) : \theta_{\omega_0 \omega_1} \cdots \theta_{\omega_{N-1} \omega_N} > 0\}$. This choice of \bar{r} will be clarified in what follows.

Now, fix $k = 1, 2, \dots, |I|$ and fix a profile $\hat{s} \in S^k$. By the definition of $\underline{V}_\omega^{\hat{s}}$ ($\omega \in \Omega$), there exists $h_\omega \in H$ such that $s(h_\omega) = \hat{s}$, $\omega(h_\omega) = \omega$, and

$$\underline{V}_\omega^{\hat{s}} > V(f|h_\omega) - \frac{\epsilon \underline{\theta}}{2|I||\Omega|}. \quad (3.6)$$

We have

$$V(f|h_\omega) = (1 - \sum_{\omega' \in \Omega} \theta_{\omega \omega'}) u(\hat{s}) + \sum_{\omega' \in \Omega} \theta_{\omega \omega'} V(f|h_\omega \circ \omega'). \quad (3.7)$$

Since f is a perfect equilibrium, for each $\omega' \in \Omega_j$ with $j \in I$,

$$V(f|h_\omega \circ \omega') \geq V(f|h_\omega \circ (\omega'; \hat{s}_j)) \geq \underline{V}_{\omega'}^{\hat{s}}, \quad (3.8)$$

where the second inequality holds by the definition of $\underline{V}_\omega^{\hat{s}}$. Similarly, if $\hat{s}_j \neq s_j^*$, then

$$V(f|h_\omega \circ \omega') \geq V(f|h_\omega \circ (\omega'; s_j^*)) \geq \underline{V}_{\omega'}^{\hat{s} \setminus s_j^*}, \quad (3.9)$$

Also, by the definition of $\underline{V}_{\omega'}^{\hat{s}}$ for $\omega \in \Omega_0$, we have

$$V(f|h_\omega \circ \omega') \geq \underline{V}_{\omega'}^{\hat{s}}, \quad \forall \omega' \in \Omega_0. \quad (3.10)$$

Substituting (3.8) and (3.10) into (3.7), we obtain

$$V(f|h_\omega) \geq (1 - \sum_{\omega' \in \Omega} \theta_{\omega \omega'}) u(\hat{s}) + \sum_{\omega' \in \Omega} \theta_{\omega \omega'} \underline{V}_{\omega'}^{\hat{s}}. \quad (3.11)$$

Inequalities (3.6) and (3.11) imply

$$\underline{V}_\omega^{\hat{s}} \geq (1 - \sum_{\omega' \in \Omega} \theta_{\omega \omega'}) u(\hat{s}) + \sum_{\omega' \in \Omega} \theta_{\omega \omega'} \underline{V}_{\omega'}^{\hat{s}} - \frac{\epsilon \underline{\theta}}{2|I||\Omega|}. \quad (3.12)$$

Also, if $s_{i(\omega^*)}(h) \neq s_{i(\omega^*)}^*$ holds for $\omega^* \in \Omega$, then

$$\underline{V}_\omega^{\hat{s}} \geq (1 - \sum_{\omega' \in \Omega} \theta_{\omega\omega'})u(\hat{s}) + \sum_{\omega' \neq \omega^*} \theta_{\omega\omega'} \underline{V}_{\omega'}^{\hat{s}} + \theta_{\omega\omega^*} \underline{V}_{\omega^*}^{s \setminus s_i^*(\omega^*)} - \frac{\epsilon\theta}{2|I||\Omega|}. \quad (3.13)$$

By definition, $\underline{V}_{\omega_0}^{\hat{s}} = \underline{V}^{\hat{s}}$ for some ω_0 . Take such ω_0 . There exists $i \in I$ such that $\hat{s}_i \neq s_i^*$. Then there exists a chain $\omega_0, \omega_1, \dots, \omega_N$ in Ω ($N < |\Omega|$) with $\omega_N \in \Omega_i$ such that $p_{\omega_{n-1}\omega_n} > 0$ for all $n = 1, \dots, N$. Sequentially substituting ω_n ($n = 0, 1, \dots, N-1$ in place of ω in (3.12), and substituting ω_N in place of ω^* in (3.13) we obtain

$$\begin{aligned} \underline{V}^{\hat{s}} &> (1 - \sum_{\omega' \in \Omega} \theta_{\omega_0\omega'})u(\hat{s}) + \sum_{\omega' \neq \omega_1} \theta_{\omega_0\omega'} \underline{V}_{\omega'}^{\hat{s}} \\ &\quad + \theta_{\omega_0\omega_1} \left[(1 - \sum_{\omega' \in \Omega} \theta_{\omega_1\omega'})u(\hat{s}) + \sum_{\omega' \neq \omega_2} \theta_{\omega_1\omega'} \underline{V}_{\omega'}^{\hat{s}} \right] \\ &\quad + \dots \\ &\quad + \theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-2}\omega_{N-1}} \left[(1 - \sum_{\omega' \in \Omega} \theta_{\omega_N\omega'})u(\hat{s}) + \sum_{\omega' \neq \omega_N} \theta_{\omega_{N-1}\omega'} \underline{V}_{\omega'}^{\hat{s}} \right] \\ &\quad + \theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-1}\omega_N} \underline{V}_{\omega_N}^{s \setminus s_i^*} - \frac{\epsilon\theta}{2|I|}. \end{aligned} \quad (3.14)$$

Using $\underline{V}_\omega^{\hat{s}} \geq \underline{V}^{\hat{s}}$ ($\forall \omega \in \Omega$), we have

$$\underline{V}^{\hat{s}} > \left[1 - \frac{\theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-1}\omega_N}}{\theta(\omega_0, \dots, \omega_N)} \right] u(\hat{s}) + \frac{\theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-1}\omega_N}}{\theta(\omega_0, \dots, \omega_N)} \underline{V}^{s \setminus s_i^*} - \frac{\epsilon}{2|I|} \quad (3.15)$$

Since \bar{r} was chosen to satisfy (3.5), if $r < \bar{r}$, then

$$\min_s u(s) > \left[1 - \frac{\theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-1}\omega_N}}{\theta(\omega_0, \dots, \omega_N)} \right] u^* + \frac{\theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-1}\omega_N}}{\theta(\omega_0, \dots, \omega_N)} \min_s u(s) - \frac{\epsilon}{2|I|} \quad (3.16)$$

Comparing (3.15) with (3.16), for any $r < \bar{r}$ then

$$\underline{V}^{\hat{s}} > \underline{V}^{s \setminus s_i^*} - \frac{\epsilon}{|I|}$$

holds where, by construction, $s \setminus s_i^* \in S^{k-1}$.

Since S^0 is the singleton set, $\{s^*\}$, we have $\underline{V}^s > u^* - \frac{\epsilon}{|I|}$ for all $s \in S^1$. It follows that $\underline{V}^s > \underline{V}^{s \setminus s_i^*} - \frac{\epsilon}{|I|}$ holds for all $s \in S^2$, and hence it holds for all $s \in S^k$. Therefore, by recursive substitution,

$$\underline{V}^{\hat{s}} > u^* - \epsilon, \quad \forall \hat{s}$$

□

4. General Coordination Games

Unfortunately, It turns out that uniqueness is not generally attainable in coordination games. In the 2×2 game G_2 below there are Perfect equilibria in which, due to asymmetric costs of miscoordination, each player receives a payoff bounded away from $u_i^* = 100$ for all r .

2			
		s_2^*	\bar{s}_2
1	s_1^*	100, 100	-200, 50
	\bar{s}_1	50, -200	10, 10
G_2			

Using the Poisson revision process of Example 2, suppose $\Delta = 1/\lambda p_1 = 1/\lambda p_2$ denoting the average delay, then for either Δ or r sufficiently small, the continuation value giving payoff (10, 10) can be supported by the strategy in which both players remain forever in \bar{s} once it is reached. If the initial profile is s^* both individuals switch to \bar{s}_i at the first revision opportunity; from profile (\bar{s}_1, s_2^*) player 2 switches to \bar{s}_2 , and from profile (s_1^*, \bar{s}_2) player 1 switches. If player 1, for example, chooses to remain in s^* at his revision opportunity, then he receives value $(1 - 2\theta)100 + \theta(50(1 - \gamma) + 10\gamma) + \theta(-200(1 - \gamma) + 10\gamma)$ where $\gamma = \lambda/(r + \lambda)$ and $\theta = \lambda/(r + 2\lambda)$. If player 1 switches, he receives $50(1 - \gamma) + 10\gamma$. It can be verified then that he strictly prefers to switch for small Δ . Since each player prefers the payoff from switching first, the profile s^* is not sustained.⁵

4.1. Generic Payoffs and Optimality

Is the optimality result nongeneric in payoff space? It seems so if, for each game, we look at the set of equilibria as the discount rate approaches zero. If, on the other hand, we fix the discount sufficiently small, then for each pure coordination game, we always find an open neighborhood in payoff space such that in every game in the neighborhood, the Pareto efficient outcome is the unique PE outcome. The following theorem establishes it.

Theorem 3. *Given $\epsilon > 0$ and any asynchronously repeated game $\langle G, \nu, (p_{\omega\omega'}), r \rangle$ in which $G = (I, S, u)$ is a pure coordination game with u^* as the unique Pareto efficient outcome, there exists $\bar{r} > 0$ such that for any $r \in (0, \bar{r})$, there exists an open dense subset $\mathcal{U} \subset \mathfrak{R}^{\times_{i \in I} S_i}$ with $u \in \mathcal{U}$ such that in an asynchronously repeated game $\langle (I, S, u'), \nu, (p_{\omega\omega'}), r \rangle$ with $u' \in \mathcal{U}$, every continuation value in any perfect equilibrium is at least $u^* - \epsilon$.*

Theorem 3 and the example that uses stage game G_2 show that order of limits matters. The example with G_2 fixes the stage game and then varies r . By contrast, the hypothesis of Theorem 3

⁵This example shows that the optimality result utilizes a different logic than that used in, say, Gale (1995) in which players' decision nodes arrive sequentially, but the game ends in a finite number of steps.

fixes r and then varies G . The former will surely be more familiar to those familiar with the Folk Theorem. The latter may be more useful if it is the discount rate r , rather than the stage payoffs, which is pinned down by exogenous data.

proof of Theorem 3. Take as given an asynchronously repeated game $\langle G, \nu, (p_{\omega\omega'}), r \rangle$ in which $G = (I, S, u)$ is a pure coordination game, and $s^* \in S$ is the unique Pareto efficient outcome. Also take $\epsilon > 0$ as given. Any affine transformation of u will give the same result in the following analysis. Analogously to the proof of Theorem 2 we set \bar{r} so as to satisfy

$$\min_{\substack{(\omega_0, \omega_1, \dots, \omega_N) \in \Omega^{N+1} \\ 2 \leq N \leq |\Omega| - 1}} \left\{ \frac{\theta_{\omega_0\omega_1} \cdots \theta_{\omega_{N-1}\omega_N}}{\theta(\omega_0, \dots, \omega_N)} : \theta_{\omega_0\omega_1} \cdots \theta_{\omega_{N-1}\omega_N} > 0 \right\} > 1 - \frac{\epsilon}{3|I|} \frac{1}{u^* - \min_s u(s)},$$

where, just as in (3.5),

$$\theta(\omega_0, \dots, \omega_N) = 1 - \sum_{\omega' \neq \omega_1} \theta_{\omega_0\omega'} - \theta_{\omega_0\omega_1} \sum_{\omega' \neq \omega_2} \theta_{\omega_1\omega'} - \cdots - \theta_{\omega_0\omega_1} \cdots \theta_{\omega_{N-2}\omega_{N-1}} \sum_{\omega' \neq \omega_N} \theta_{\omega_{N-1}\omega'}.$$

Such an \bar{r} can be found since both $\theta_{\omega_0\omega_1} \cdots \theta_{\omega_{N-1}\omega_N}$ and θ converge to $p_{\omega_0\omega_1} \cdots p_{\omega_{N-1}\omega_N} > 0$ as \bar{r} goes to zero. Fix $r \in (0, \bar{r})$. Again let $\underline{\theta} = \min \theta(\omega_0, \dots, \omega_N)$.

Consider a neighborhood of u given by $\mathcal{U} = \{u' \in \mathfrak{R}^S \mid \forall s \in S, |u'_i(s) - u_i(s)| < \eta/4\}$ where η satisfies

$$0 < \eta < \frac{1}{2} \min \left\{ u^* - u^{**}, \frac{\epsilon}{2|I||\Omega|}, \frac{\epsilon}{3(|I| + 1)} \left[1 - \int_0^\infty e^{-r\tau} d\nu(\tau) \right] \right\},$$

where $u^{**} = \max_{s \neq s^*} u(s)$, i.e., the second highest payoff. In the following, we consider an asynchronously repeated game $\Gamma' = \langle (I, S, u'), \nu, (p_{\omega\omega'}), r \rangle$ with $u' \in \mathcal{U}$.

Now, fix a perfect equilibrium f of Γ' . For each $i \in I$, each $s \in S$, and each $\omega \in \Omega$, we let $\underline{V}_{i\omega}^s = \inf_{s(h)=s, \omega(h)=\omega} V_i(f|h)$, and $\underline{V}_i^s = \inf_{s(h)=s} V_i(f|h)$. Note that $\underline{V}_i^s = \min_{\omega \in \Omega} \underline{V}_{i\omega}^s$.

We first show the counterpart of Theorem 1. For any $\delta > 0$ there exists $h \in H$ such that $s(h) = s^*$ and

$$\underline{V}_i^{s^*} > V_i(f|h) - \delta, \quad \forall i \in I. \quad (4.1)$$

The continuation value for player $i \in I$ after h is given by

$$V_i(f|h) = \left(1 - \sum_{\omega \in \Omega} \theta_{\omega(h)\omega} \right) u'_i(s^*) + \sum_{\omega \in \Omega} \theta_{\omega(h)\omega} V_i(f|h \circ \omega). \quad (4.2)$$

If $\omega(h) \in \Omega_0$, then $s(h \circ \omega) = s^*$ and therefore, $V_i(f|h \circ \omega) \geq \underline{V}_i^{s^*}$. Observe also that since f is a perfect equilibrium strategy profile, it must be the case that for all $\omega \in \Omega_j$ ($j \in I$),

$$V_j(f|h \circ \omega) \geq V_j(f|h \circ (s_j^*; \omega)) \geq \underline{V}_j^{s_j^*} \quad (4.3)$$

where the second inequality holds due to the definition of $\underline{V}_j^{s_j^*}$. Since $|u(s) - u_i(s)| < \eta/4$, $|V_j(f|h) - V_i(f|h)| < \eta/2$ holds for all $i, j \in I$, all f and all $h \in H$, and therefore, (4.3) implies

$$V_i(f|h \circ \omega) \geq \underline{V}_i^{s^*} - \eta. \quad (4.4)$$

Substituting (4.4) into (4.2) and using (4.1), we obtain

$$\underline{V}_i^{s^*} > (1 - \sum_{\omega' \in \Omega} \theta_{\omega(h)\omega'}) u'_i(s^*) + \sum_{\omega' \in \Omega} \theta_{\omega(h)\omega'} \underline{V}_i^{s^*} - \delta - \eta \quad (4.5)$$

Inequality (4.5) implies

$$\underline{V}_i^{s^*} > u'_i(s^*) - \frac{\delta + \eta}{1 - \int_0^\infty e^{-r\tau} d\nu(\tau)}.$$

Since δ is arbitrary and independent of other variables, the definition of η implies $\underline{V}_i^{s^*} > u^* - \epsilon/(|I| + 1)$.

This second part mirrors the proof of Theorem 2. We will show that for all $k = 1, 2, \dots, |I|$, after h with $s(h) \in S^k$, its continuation value satisfies

$$V_i(f|h) > \underline{V}_i^{s'} - \frac{\epsilon}{|I| + 1}, \quad (4.6)$$

for some $s' \in S^{k-1}$ if $r < \bar{r}$. Backward induction implies that (4.6) holds for all $s' \in S^{k-1}$ if $r < \bar{r}$. Once we show (4.6) for all s' and k 's, we verify that $0 < r < \bar{r}$ implies that for all history $h \in H$, $V_i(f|h) > u'_i(s^*) - \epsilon$. Moreover, recall that the choice of \bar{r} is independent of f . Thus, the proof will be completed.

Fix $k = 1, 2, \dots, |I|$ and $\hat{s} \in S^k$. By the definition of $\underline{V}_{i\omega}^{\hat{s}}$ ($\omega \in \Omega$), there exists $h_\omega \in H$ such that $s(h_\omega) = \hat{s}$, $\omega(h_\omega) = \omega$, and

$$\underline{V}_{i\omega}^{\hat{s}} > V_i(f|h_\omega) - \frac{\epsilon\theta}{3(|I| + 1)|\Omega|}. \quad (4.7)$$

We have

$$V_i(f|h_\omega) = (1 - \sum_{\omega' \in \Omega} \theta_{\omega\omega'}) u'_i(\hat{s}) + \sum_{\omega' \in \Omega} \theta_{\omega\omega'} V_i(f|h_\omega \circ \omega'). \quad (4.8)$$

Since f is a perfect equilibrium, for each $\omega' \in \Omega_j$ with $j \in I$,

$$V_j(f|h_\omega \circ \omega') \geq V_j(f|h_\omega \circ (\hat{s}_j; \omega)) \geq \underline{V}_{j\omega}^{\hat{s}}, \quad (4.9)$$

where the second inequality holds by the definition of $\underline{V}_{j\omega}^{\hat{s}}$. Similarly, if $\hat{s}_j \neq s_j^*$, then

$$V_j(f|h_\omega \circ \omega') \geq V_j(f|h_\omega \circ (s_j^*; \omega)) \geq \underline{V}_{j\omega}^{\hat{s} \setminus s_j^*}, \quad (4.10)$$

Also, by the definition of $\underline{V}_{i\omega'}^{\hat{s}}$ for each $\omega \in \Omega_0$, we have

$$V_i(f|h_\omega \circ \omega') \geq \underline{V}_{i\omega'}^{\hat{s}}, \quad \forall \omega' \in \Omega_0. \quad (4.11)$$

Since $|u(s) - u_i(s)| < \eta/4$, then $|V_j(f|h) - V_i(f|h)| < \eta/2$ holds for all $i, j \in I$, all f , all s , and all $h \in H$, and therefore, (4.9) and (4.10) imply

$$V_i(f|h_\omega \circ \omega') \geq \underline{V}_{i\omega'}^{\hat{s}} - \eta, \quad (4.12)$$

and

$$V_i(f|h_\omega \circ \omega') \geq \underline{V}_{i\omega'}^{\hat{s} \setminus s_j^*} - \eta, \quad (4.13)$$

respectively. Substituting these inequalities into (4.8), we obtain

$$V_i(f|h_\omega) \geq (1 - \sum_{\omega' \in \Omega} \theta_{\omega\omega'})u'_i(\hat{s}) + \sum_{\omega' \in \Omega} \theta_{\omega\omega'}V_{i\omega'}^{\hat{s}} - \eta. \quad (4.14)$$

Inequalities (4.7) and (4.14) imply

$$\underline{V}_{i\omega}^{\hat{s}} \geq (1 - \sum_{\omega' \in \Omega} \theta_{\omega\omega'})u'_i(\hat{s}) + \sum_{\omega' \in \Omega} \theta_{\omega\omega'}V_{i\omega'}^{\hat{s}} - \frac{\epsilon\theta}{3(|I|+1)|\Omega|} - \eta. \quad (4.15)$$

Since i was arbitrarily chosen, (4.15) holds for all $i \in I$. By definition, $\underline{V}_{i\omega_0}^{\hat{s}} = \underline{V}_i^{\hat{s}}$ for some ω_0 . Take such ω_0 . There exists $i \in I$ such that $\hat{s}_i \neq s_i^*$. Then there exists a chain $\omega_0, \omega_1, \dots, \omega_N$ with $N < |\Omega|$ and $\omega_N \in \Omega_i$ such that $p_{\omega_{n-1}\omega_n} > 0$ for all $n = 1, \dots, N$. Sequentially substituting ω_n ($n = 0, 1, \dots, N$ in place of ω in (4.15) and applying (4.13), we obtain

$$\begin{aligned} \underline{V}_i^{\hat{s}} &\geq (1 - \sum_{\omega' \in \Omega} \theta_{\omega_0\omega'})u'_i(\hat{s}) + \sum_{\omega' \neq \omega_1} \theta_{\omega_0\omega'}V_{i\omega'}^{\hat{s}} \\ &\quad + \theta_{\omega_0\omega_1} \left[(1 - \sum_{\omega' \in \Omega} \theta_{\omega_1\omega'})u'_i(\hat{s}) + \sum_{\omega' \neq \omega_2} \theta_{\omega_1\omega'}V_{i\omega'}^{\hat{s}} \right] \\ &\quad + \dots \\ &\quad + \theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-2}\omega_{N-1}} \left[(1 - \sum_{\omega' \in \Omega} \theta_{\omega_N\omega'})u'_i(\hat{s}) + \sum_{\omega' \neq \omega_N} \theta_{\omega_{N-1}\omega'}V_{i\omega'}^{\hat{s}} \right] \\ &\quad + \theta_{\omega_0\omega_1} \theta_{\omega_{N-1}\omega_N} \underline{V}_{i\omega_N}^{\hat{s} \setminus s_i^*} - \frac{\epsilon\theta}{3(|I|+1)} - \eta|\Omega|. \end{aligned} \quad (4.16)$$

Using $\underline{V}_{i\omega}^{\hat{s}} \geq \underline{V}_i^{\hat{s}}$ ($\forall \omega \in \Omega$), we have

$$\underline{V}_i^{\hat{s}} \geq \left[1 - \frac{\theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-1}\omega_N}}{\theta(\omega_0, \dots, \omega_N)} \right] u'_i(\hat{s}) + \frac{\theta_{\omega_0\omega_1} \dots \theta_{\omega_{N-1}\omega_N}}{\theta(\omega_0, \dots, \omega_N)} \underline{V}_i^{\hat{s} \setminus s_i^*} - \frac{2\epsilon}{3(|I|+1)}$$

Thus, for all $r < \bar{r}$,

$$\underline{V}_i^{\hat{s}} > \underline{V}_i^{\hat{s} \setminus s_i^*} - \frac{\epsilon}{|I|+1},$$

where, by construction, $\hat{s} \setminus s_i^* \in S^{k-1}$. □

4.2. More on Genericity

Though uniqueness of the Pareto optimal outcome is not generally possible, i.e., not possible for all discount rates, for general coordination games with asymmetric costs of miscoordination we do establish asymptotic uniqueness for an interval of discount rates. This stands in contrast with standard repeated games. Pareto inferior Nash equilibria of the stage game are always equilibria of the standard repeated game, regardless of the value of the discount rate. We show that the inferior Nash equilibrium is not sustainable in a neighborhood of discount rates for the class of symmetric 2×2 coordination games given in G_3 .

2

		s_2^*	\bar{s}_2
1	s_1^*	u^*, u^*	$\beta, 0$
	\bar{s}_1	$0, \beta$	$1, 1$
G_3			

Theorem 4. Consider any asynchronously repeated game $\Gamma = \langle G_3, \nu, (p_{\omega\omega'}), r \rangle$ with G_3 as above where $u^* > 2$ and $\beta > 0$. Then there exists a nonempty open set $R \in (0, \infty)$ such that if $r \in R$ then for any perfect equilibrium f ,

$$V_i(f|h) = u^*, \forall i, \forall h \text{ with } s(h) = s^*,$$

and with probability one, there is some history h with $s(h) = s^*$ is reached.

Note that from the proof it will be clear that the parametric part of G_3 is not essential to the argument.

proof of Theorem 4. Fix a perfect equilibrium f . First, let $\underline{V}_i = \inf_{s(h)=s^*} V_i(f|h)$ ($i = 1, 2$). Assume the contrary, i.e., that $\underline{V}_1 < u^*$. Note that $\underline{V}_2 < u^*$ holds, too. By the recursive formula for V and the above contrapositive assumption, for all $\epsilon > 0$ there exists $h = h_\epsilon \in H$ such that player 2 switches to \bar{s}_2 at $h' = h \circ \omega'$ with positive probability where $p_{\omega(h)\omega'} > 0$, and

$$\underline{V}_1 > V_1(f|h) - \epsilon$$

and

$$\underline{V}_1 > V_1(f|h \circ (\omega'; \bar{s}_2)) \tag{4.17}$$

hold. By perfection of f ,

$$V_2(f|h \circ (\omega'; \bar{s}_2)) \geq V_2(f|h \circ (\omega'; s_2^*)) \geq \underline{V}_2. \tag{4.18}$$

Let $\Theta(i|\omega')$ ($i = 1, 2$) be given by

$$\begin{aligned} \Theta(i|\omega') &= \sum_{(\omega_0, \omega_1, \dots, \omega_N) \in \bar{\Omega}(i|\omega')} \theta_{\omega_0\omega_1} \cdots \theta_{\omega_{N-1}\omega_N} \\ &= \sum_{(\omega_0, \omega_1, \dots, \omega_N) \in \bar{\Omega}(i|\omega')} p_{\omega_0\omega_1} \cdots p_{\omega_{N-1}\omega_N} \left[\int_0^\infty e^{-r\tau} d\nu(\tau) \right]^N \end{aligned}$$

where

$$\bar{\Omega}(i|\omega') = \{(\omega_0, \dots, \omega_N) \in \Omega \mid \omega_0 = \omega', \forall n < N : \omega_n \notin \Omega_i, \omega_N \in \Omega_i\}.$$

That is, $\bar{\Omega}(i|\omega')$ is the set of sequences of ω 's which start with ω' and reach a state in Ω_i after N opportunities for the first time. Similarly, $\Theta(ij|\omega')$ ($i \neq j$) is given by

$$\Theta(ij|\omega') = \sum_{(\omega_0, \omega_1, \dots, \omega_N) \in \bar{\Omega}(ij|\omega')} \theta_{\omega_0 \omega_1} \cdots \theta_{\omega_{N-1} \omega_N}$$

where

$$\bar{\Omega}(ij|\omega') = \{(\omega_0, \dots, \omega_N) \in \Omega \mid \omega_0 = \omega', \exists n : \omega_n \in \Omega_i, \forall n < N : \omega_n \notin \Omega_j, \omega_N \in \Omega_j\}.$$

That is, $\bar{\Omega}(ij|\omega')$ is the set of sequences of ω 's similar to $\bar{\Omega}(j|\omega')$ but player i gets at least one revision opportunity before j .

Observe that we can write any continuation $V_i(f|h)$ as a weighted sum

$$A_h u^* + B_h u_i(s_1^*, \bar{s}_2) + C_h \cdot 1 + D_h u_i(\bar{s}_1, s_2^*)$$

with $A_h + B_h + C_h + D_h = 1$. Let h' satisfy $s(h') = (s_1^*, \bar{s}_2)$ and such that \bar{s} is assumed to be reached from h' . Observe that

$$V_1(f|h') - V_2(f|h') = \beta(B_{h'} - D_{h'})$$

Next, observe that $B_{h'} > 1 - \Theta(1|\omega)$ as the total fraction of time spent in (s_1^*, \bar{s}_2) must be at least the discounted probability that player 1 does not have a revision opportunity since \bar{s} is assumed to be reached from h' . Also, $D_{h'} < \Theta(12|\omega)$ since $\Theta(12|\omega)$ is the fastest time it takes for the players to reach (\bar{s}_1, s_2^*) and stay there permanently. Since the players will not remain in (\bar{s}_1, s_2^*) permanently under f , $D_{h'} < \Theta(12|\omega)$. Hence, the following inequality holds:

$$V_1(f|h \circ (\omega'; \bar{s}_2)) - V_2(f|h \circ (\omega'; \bar{s}_2)) \geq \beta(1 - \Theta(1|\omega') - \Theta(12|\omega')). \quad (4.19)$$

Observe that the right hand side (RHS) of (4.19) tends to be positive as r goes to infinity, and tends to be negative as r approaches to zero. Let $\underline{r}^1 > 0$ be such that the RHS of (4.19) is zero. Such an \underline{r}^1 exists since the RHS of (4.19) is continuous. Then (4.17) through (4.19) imply that for $r > \underline{r}^1$,

$$\underline{V}_1 > \underline{V}_2.$$

Switching 1 with 2, we can make the same argument to obtain

$$\underline{V}_2 > \underline{V}_1$$

for $r > \underline{r}^2$ where \underline{r}^2 is similarly defined. These two inequalities contradict with each other if $r > \max\{\underline{r}^1, \underline{r}^2\}$. Therefore, for this r , $\underline{V}_i = u^*$ for $i = 1, 2$.

Next, from an action profile (s_1^*, \bar{s}_2) , if player 2's revision node is reached, he changes his action to s_2^* . Therefore, from this profile player 1 obtains at least

$$\beta(1 - \Theta(2|\omega)) + \Theta(2|\omega)u^* \quad (4.20)$$

provided that the current state is $\omega \in \Omega$. Likewise, player 2 obtains at least

$$\beta(1 - \Theta(1|\omega)) + \Theta(1|\omega)u^* \quad (4.21)$$

from (\bar{s}_1, s_2^*) at state ω .

Now, we must show that for some $r > \max\{\underline{r}^1, \underline{r}^2\}$, at least one of the players, say player i , switches to s_i^* from profile (\bar{s}_1, \bar{s}_2) and at *some* state $\omega \in \Omega_i$. Suppose $\underline{r}^1 \geq \underline{r}^2$. The other case is analyzed in the same way. Since $\Theta(1|\omega) > \Theta(12|\omega)$, (4.19) implies that there exists $\hat{r} > \underline{r}^1$ under which $\Theta(1|\omega) > 1/2$. Let $R = (\max\{\underline{r}^1, \underline{r}^2\}, \hat{r}) \subset \mathfrak{R}$. Then for all r in R , (4.21) is greater than one since $u^* \geq 2$. It implies that staying away from s^* cannot be a perfect equilibrium. Thus, s^* is reached with probability one. But once it is reached, the action profile never leaves it if $r > \max\{\underline{r}^1, \underline{r}^2\}$. Therefore, s^* becomes the unique outcome in the long run. \square

5. A Continuity Result

Finally, we establish an ‘‘approximation Theorem’’ stating that the set of equilibrium payoffs for a one parameter family of renewal games with pure coordination stage games varies ‘‘continuously’’ between the Folk Theorem payoffs and $\{u^*\}$ as the approximating parameter varies between an asynchronously repeated game and a standard repeated game. The result demonstrates that there is no discontinuity between the Folk Theorem and ‘‘anti-Folk Theorem’’ results.

Consider a renewal game $\langle G, \nu, p_{\omega\omega'}, r \rangle$ in which G is a pure coordination game and $\Omega = 2^I$. Now define a ‘‘ δ -approximation’’ to a standard repeated game as a tuple

$$\Gamma^\delta = \langle G, \nu^\delta, p^\delta, r \rangle$$

where

$$\nu^\delta(X_k \in [1 - \delta, 1 + \delta]) \geq 1 - \delta, \forall k. \quad (5.1)$$

and

$$\sum_{j \in I} p_{C\{j\}}^\delta = \delta \text{ and } p_{CI}^\delta = 1 - \delta, \forall C \subseteq I. \quad (5.2)$$

Here, there is probability $1 - \delta$ that all players move at once at revision opportunities at the jumps of close to unit length. If $\delta = 1$ then p places full support on the singleton sets and so (a.s.) only one player moves at a time.

Given G let

$$\bar{u}^* = u(s^*)$$

$$\underline{u}^* = \min\{u(\sigma) : \sigma \text{ is a mixed Nash eqm profile}\}$$

and let

$$E(\Gamma) = E(\langle G, \nu, p, r \rangle) = \left\{ \bar{V} \in \mathfrak{R}^I \mid \exists \text{ Perfect equilibrium } f \text{ and } \exists h \in H^0 \text{ s.t. } \bar{V}_i = V_i(f|h), \forall i \right\}.$$

The set $E(\Gamma)$ denotes the Perfect equilibrium payoffs of Γ . Now let

$$\bar{E}(\Gamma) = \lim_{r \rightarrow 0} E(\Gamma) \text{ in Housdorff metric}$$

The set $\bar{E}(\Gamma)$ is the limit of the set of Perfect equilibrium payoffs of Γ taken as r approaches zero.

Theorem 5. Given a renewal game Γ^δ with pure coordination, there is a $\eta(\delta) > 0$ such that

(i) $\bar{E}(\Gamma^\delta) = [\bar{u}^* - \eta(\delta), \bar{u}^*]$.

(ii) $\eta(\cdot)$ is continuous, and $\eta(\delta) \rightarrow 0$ as $\delta \rightarrow 1$ while $\eta(\delta) \rightarrow \bar{u}^* - \underline{u}^*$ as $\delta \rightarrow 0$.

Before proceeding with the proof we should emphasize that the result does *not* prove that the standard repeated model is robust since the δ approximation is just one way of defining the standard model as a limit of sequences of renewal games. Recall Example 3 in Section 2 which provides an alternative definition. In that formulation, the standard model is not robust since it is the limit of sequences of asynchronously repeated games defined by Example 3 (as opposed to being a limit of sequences of games in which *some* decisions are made asynchronously — which is the case here).

proof of Theorem 5. Fix $\epsilon > 0$. Fix $\delta > 0$, and $r > 0$, and an equilibrium f of Γ^δ given r . The argument mirrors that of Theorems 1 and 2. We first give a lower bound to PE payoffs, then show that the payoffs that exceed this bound less ϵ are achieved if r is sufficiently small. This lower bound will be shown to be $\bar{u}^* - \eta(\delta)$ with $\eta(\delta)$ satisfying the requisite properties.

Given the δ approximation, Γ^δ , define (as in Theorem 1) $\underline{V} = \inf\{V(f|h) : s(h) = s^*\}$. Then for any $\epsilon > 0$ there is some history h with $s(h) = s^*$ such that

$$\underline{V} > [1 - \sum_{\omega' \in \Omega} \theta_{\omega(h)\omega'}] \bar{u}^* + \sum_{\omega' \in \Omega} \theta_{\omega(h)\omega'} V(f|h \circ \omega') - \epsilon \quad (5.3)$$

Given the construction of the δ approximation we can rewrite the right hand side of (5.3) as

$$[1 - \sum_{\omega' \in \Omega} \theta_{\omega(h)\omega'}] \bar{u}^* + \int e^{-rt} d\nu \left(\sum_{i \in I} p_{\omega(h)\{i\}}^\delta V(f|h \circ \{i\}) + (1 - \delta) V(f|h \circ I) \right) - \epsilon$$

With Perfection it follows that $V(f|h \circ \{i\}) \geq V(f_{-i}, \hat{f}_i|h \circ \{i\})$ for all \hat{f}_i . In particular this inequality holds for any \hat{f}_i with $\hat{f}_i(h \circ \{i\}) = s_i^*$. Therefore, we have

$$\underline{V} > \frac{1 - \int e^{-rt} d\nu}{1 - \delta \int e^{-rt} d\nu} \bar{u}^* + \frac{(1 - \delta) \int e^{-rt} d\nu V(f|h \circ I) - \epsilon}{1 - \delta \int e^{-rt} d\nu} \quad (5.4)$$

If individuals use their minimax strategies in state $h \circ I$ then (5.4) becomes

$$\underline{V} > \frac{1 - \int e^{-rt} d\nu}{1 - \delta \int e^{-rt} d\nu} \bar{u}^* + \frac{(1 - \delta) \int e^{-rt} d\nu \underline{u}^* - \epsilon}{1 - \delta \int e^{-rt} d\nu} \quad (5.5)$$

Since (5.5) holds for all $\epsilon > 0$, there is a number $\eta(\delta) > 0$ which satisfies the limit properties of (ii). It is given by

$$\eta(\delta) = \lim_{\epsilon \rightarrow 0} \left[\frac{(1 - \delta) \int e^{-rt} d\nu}{1 - \delta \int e^{-rt} d\nu} \bar{u}^* - \frac{(1 - \delta) \int e^{-rt} d\nu \underline{u}^* - \epsilon}{1 - \delta \int e^{-rt} d\nu} \right] \quad (5.6)$$

Observe that $\eta(\cdot)$ is continuous. Now the same backward induction argument as in Theorem 2 shows that there is some $\bar{r} > 0$ such that if $0 < r \leq \bar{r}$ then $V(f|h)$ is greater than $\bar{u}^* - \eta(\delta) - \epsilon$ given any initial history h .

It remains to show that any payoff which exceeds this lower bound can be achieved as a limit payoff of a sequence of Perfect equilibrium payoffs as $r \rightarrow 0$. Let (\hat{u}_1, \hat{u}_2) denote a payoff that exceeds the lower bound in (5.5). We construct the equilibrium f to approximate (\hat{u}_1, \hat{u}_2) if r is small. For any history h and at any state $\omega = \{i\}$, $i \in I$, let $f_i(h \circ \{i\}) = s_i^*$. That is, at all asynchronous decision states, play remains in or returns to s^* as soon as possible. If h is on the equilibrium path, we define at decision state $\omega = I$, $f_i(h^t \circ I) = \bar{\sigma}_i$ for each i where $\bar{u}_i = \sum_{s \in S} \bar{\sigma}_i(s) u_i(s)$ is defined implicitly by

$$\hat{u}_i = \frac{1 - \int e^{-rt} d\nu}{1 - \delta \int e^{-rt} d\nu} \bar{u}_i + \frac{(1 - \delta) \int e^{-rt} d\nu}{1 - \delta \int e^{-rt} d\nu} \bar{u}_i \quad (5.7)$$

Since there is a mixed strategy that can achieve any symmetric payoff in $[\underline{u}^*, \bar{u}^*]$ such a $\bar{\sigma}$ and \bar{u} can be found. If there is ever a deviation from this strategy we let $f_i(h^t \circ I) = \underline{\sigma}_i$ after the deviation history h^t where $\underline{\sigma}$ achieves the minimax payoff \underline{u}^* . Since $\hat{u}_i > \bar{u}_i - \eta(\delta)$, the minimax punishment deters deviations from the prescribed f . Also, since $\bar{u}_i - \eta(\delta)$ is itself a Perfect equilibrium continuation payoff, f is a Perfect equilibrium which achieves payoff profile \hat{u} . \square

6. Related Literature

Asynchronous choice models have been studied to some extent in various settings. As a points of comparison with related models, we emphasize the work on bargaining by Perry and Reny (1993), the dynamic oligopoly problems studied by Maskin and Tirole (1988), Farrell and Saloner (1985), Gale (1995), and Morris (1995), and our previous work (some with Rafael Rob), on asynchronously repeated games with evolutive dynamics (Matsui and Rob (1992), Lagunoff and Matsui (1995), and Lagunoff (1995)).

In the Perry/Reny model, individuals can move at any time, with a small delay after each player moves. They show that if players can react to offers instantaneously then the set of equilibria are bounded by the first and second mover payoffs, resp., of the Rubinstein bargaining model. The Perry/Reny model offers the advantage over our framework that the choice of timing of moves is endogenous subject to a fixed waiting time that must be endured after a player makes an offer. However, since players have the option of moving simultaneously, it appears that in the adaptation of their model to the repeated game, each of the static Nash equilibria of a coordination game will persist as Perfect equilibria.

Maskin and Tirole rule out simultaneous moves by considering an alternating move game in which, in one variant of the model, each firm sets capacity given the (temporarily) fixed capacity of rivals. The timing structure of the model is subsumed in our general definition of asynchronously repeated games, however, their interest is with strategic behavior that depends only on payoff relevant information, i.e., natural state variables.

Works by Farrell and Saloner (1985), Gale (1995), and Morris (1995) are also related to the present paper. Farrell/Saloner and Gale show that in an n -person coordination game with asynchronous choice and finitely many moves, a Pareto efficient outcome is attained if players are sufficiently patient. Backward induction is used to establish that when, say, first $k < n$ players cooperate, the $(k + 1)$ th player also cooperates. Though its logic looks similar to ours at a first glance, it is very different. The difference is manifested in Game G_2 . In this game, s^* is uniquely attained in the framework of Farrell/Saloner and Gale, while \bar{s} is also reached infinitely often in our framework. The difference is that there is no fear of “retaliation” for the last person in theirs, while there is always a chance of “retaliation” in our framework. When the game played is a pure coordination game, this “retaliation” does not work since one who “retaliates” also harms himself to the same degree.

Morris considers a strategic environment similar to Farrell/Saloner and Gale except that the entry times into the game of each of the n individuals are not common knowledge. Each player observes his own “clock” but knows only the support of the distribution over others’ clocks. In contrast to the first two, Morris shows that optimal coordination is, in fact, never achieved. As with the other two, the nonrepetitive strategic environment prevents “retaliation.” Without this threat, synchronized coordination unravels when the timing of moves is not common knowledge.

More closely related to the present paper are the Matsui/Rob, Lagunoff/Matsui, and Lagunoff evolutive models which assume the Poisson revision process of Example 2 in Section 2. New individuals were assumed to enter the game stochastically and asynchronously. In the first two models heterogeneous forecasts across generations provided the impetus for change. In the last model, perfect foresight was assumed and, for this reason, is closest to the present equilibrium model.⁶

There is also a sizable literature which characterizes equilibria of repeated games that differ in other respects from the standard model. Two distinct (though related) approaches may be found. In one, restrictions are placed on repeated game equilibria by considering those that are limit points of sequences of equilibria of perturbed games with perturbations in the information structure. Often these perturbations allow the players to precommit to particular outcomes through reputation or informational leakage. Examples include Fudenberg and Levine (1987), Matsui (1989), and Aumann and Sorin (1989). All of these have unique “limit equilibria” for a certain class of stage games.

Fudenberg and Levine (FL) (1987), for instance, characterize outcomes of a repeated game in which a long run player faces a sequence of short run or “myopic” players. They show that only payoffs that dominate a certain “stackelberg” payoff arise when the short run players have a small amount of uncertainty regarding the long run player’s “type.”⁷ However, since FL are primarily

⁶Related work includes Blume (1993), Matsui (1994), and Matsui and Matsuyama (1995). Individuals’ decisions in these papers are asynchronous, however, the players are either assumed to be myopic, or are randomly matched in a large population so that no player accounts for the intertemporal effects of his own behavior.

⁷The “stackelberg” payoffs are those that arise if the long run player could commit to a behavior

interested in modeling reputation, they consider only “limit equilibria” as the uncertainty becomes negligible. This, together with the presence of myopic players on one side of the game eliminates some of multiplicity created by infinite repetition.

A second approach uses bounds on strategic complexity to restrict equilibria in repeated games. Examples are Rubinstein (1986), Abreu and Rubinstein (1988), Kalai, Samet, and Stanford (1988), and Cho and Li (1995). In some of these, e.g., Rubinstein (1986), stage game Nash equilibria may be eliminated as solutions even if complexity costs enter into preferences lexicographically.⁸

Both of the above approaches tend to look at a more general class of stage games than is considered in the present work. As with standard repeated games, we differ from these by considering fully rational players (subject to the simplifying recursivity assumption), and we consider all the Perfect equilibria of the repeated game.

It may be worthwhile to look at asynchronous choice with strategies of bounded complexity. One useful starting point is to generalize Maskin and Tirole’s state variable approach to other repeated renewal games. In our terminology, the “natural state space” is (Ω, S) which defines who can physically move at a decision point.

References

- [1] Abreu, D.: On the theory of infinitely repeated games with discounting. *Econometrica*, 56: 383-398 (1988).
- [2] Abreu, D., Rubinstein, A: The structure of Nash equilibrium in repeated games with finite automata. *Econometrica*, 56: 1259-1281.
- [3] Aumann, R.: Survey of repeated games. In *Essays in Game Theory and Mathematical Economics in Honor of Oskar Morgenstern*, Mannheim: Bibliographisches Institut (1981).
- [4] Aumann, R., Sorin, S.: Cooperation and bounded recall. *Games Econ. Behavior*, 1: 5-39 (1989).
- [5] Blume, L.: The statistical mechanics of strategic interaction. *Games Econ. Behavior*, 5: 387-424 (1993).
- [6] Cho, I.-K., Li, H.: Complexity and neural networks. Mimeo, University of Chicago (1995).
- [7] Farrell, J., Saloner, G.: Standardization, compatibility, and innovation. *Rand J. Econ.*, 16: 70-83 (1985).

strategy taking as given the myopic best response of the short run players. It coincides with the Pareto dominant payoff in pure coordination games.

⁸Without these costs it is not true necessarily that the structure of equilibrium payoffs differ fundamentally from the Folk Theorem. See Kalai and Stanford (1988).

- [8] Fudenberg, D., Levine, D.K.: Reputation and equilibrium selection in games with a patient player. *Econometrica*, 57: 759-78 (1989).
- [9] Fudenberg, D., Maskin, E.: The Folk Theorem in repeated games with discounting or with incomplete information. *Econometrica*, 54: 533-56 (1986).
- [10] Gale, D.: Dynamic coordination games. *Econ. Theory*, 5: 1-18 (1995).
- [11] Kalai, E., Samet, D., Stanford, W.: Reactive equilibria. *International J. Game Theory*, 176-86 (1988).
- [12] Kalai, E., Stanford, W.: Finite rationality and interpersonal complexity in repeated games. *Econometrica*, 56: 397-410 (1988).
- [13] Lagunoff, R.: On the evolution of pareto optimal behavior in repeated coordination problems. *Mimeo, University of Pennsylvania* (1995).
- [14] Lagunoff, R., Matsui, A.: Evolution in mechanisms for public projects. *Econ. Theory*, 6: 195-223 (1995).
- [15] Maskin, E., Tirole, J.: A theory of dynamic oligopoly, I: Overview and quantity competition with large fixed costs. *Econometrica*, 56: 549-70 (1988).
- [16] Matsui, A.: Information leakage forces cooperation. *Games Econ. Behavior*, 1: 94-115 (1989).
- [17] Matsui, A.: Evolution and rationalizability. *Mimeo, University of Pennsylvania, March* (1993).
- [18] Matsui, A., Matsuyama, K.: An approach to equilibrium selection. *J. Econ. Theory*, 65: 415-34 (1995).
- [19] Matsui, A., Rob, R.: Evolution with forward looking players. *Mimeo, University of Pennsylvania* (1992).
- [20] Morris, S.: Cooperation and timing. *Mimeo, University of Pennsylvania* (1995).
- [21] Perry, M., Reny, P.: A non-cooperative bargaining model with strategically timed offers. *J. Econ. Theory*, 59: 50-77 (1993).
- [22] Rubinstein, A.: Finite autonomous play the repeated prisoner's dilemma. *J. Econ. Theory*, 39: 83-96 (1986).

This is a list of recent CARESS Working Papers. A complete list (dating from inception of the series) can be obtained by writing to:

Mrs. Kelly A. Quinn
CARESS
3718 Locust Walk
McNeil Building
Philadelphia, PA 19104-6297

- 93-01 "Perpetual Randomness in Evolutionary Economics" by George J. Mailath
93-02 "Noisy Bayes Updating and the Value of Information" by Stephen Morris and Hyun Song Shin
93-03 "Buyers and Sellers" by Kenneth Burdett, Melvyn Coles, Nobuhiro Kiyotaki and Randall Wright
93-04 "AUSI Expected Utility: An Anticipated Utility Theory of Relative Disappointment Aversion" by Simon Grant and Atsushi Kajii
93-05 "Bandwagon Effects and Long Run Technology Choice" by Michihiro Kandori and Rafael Rob
93-06 "A Note on the Incentives to Agglomerate Under Demand Uncertainty" by Rafael Rob
93-07 "A Primer on Functional Differential Equations with Applications to Economics" by Paul J. Zack
93-08 "An Endogenous Rate of Time Preference, the Penrose Effect, and Dynamic Optimality of Environmental Quality" by Hirofumi Uzawa
93-09 "Bid Ask Spreads with Two Sided Private Information" by Stephen Morris
93-10 "Risk, Uncertainty and Hidden Information" by Stephen Morris
93-11 "Risk Dominance and Stochastic Potential " by Stephen Morris, Rafael Rob and Hyun Song Shin
Revised and final version appears as "p-Dominance and Belief Potential," in Econometrica, 63, (January 1995), pp. 145-157.
93-12 "Structural Indifference in Normal Form Games" by George J. Mailath, Larry Samuelson and Jeroen M. Swinkels
93-13 "Search, Bargaining, Money and Prices" by Alberto Trejos and Randall Wright
93-14 "Agitators and Free Riders on the Path to Cooperation: The Evolution of Cooperation in Mechanisms for Public Projects" by Roger D. Lagunoff and Akihiko Matsui
93-15 "Depth of Knowledge and the Effect of Higher Order Uncertainty" by Stephen Morris, Andrew Postlewaite and Hyun Song Shin

Revised and final version is forthcoming in Economic Theory.

93-16 "Monetary Exchange Under Private Information: A Note" by Alberto Trejos

93-17 "Relative Income Concerns and the Rise in Married Women's Employment" by David Neumark and Andrew Postlewaite

93-18R "A Cardinal Characterization of the Rubinstein-Safra-Thomson Axiomatic Bargaining Theory" by Simon Grant and Atsushi Kajii

93-19 "Evolution and Rationalizability" by Akihiko Matsui

93-20 "Strictness, Evolutionary Stability and Repeated Games with Common Interest" by Dieter Balkenborg

93-21 "On Generic Pareto Improvement in Competitive Economies with Incomplete Asset Structures" by Alessandro Citanna and Antonio Villanacci

*93-22 "Search, Evolution and Money" Randall Wright

Revised and final version forthcoming in the Journal of Economic Dynamics and Control

*93-23 "A Note on Sunspot Equilibria in Search Models of Fiat Money" by Randall Wright

Revised and final version forthcoming in the Journal of Economic Theory

93-24 "Demand and Supply in New Markets: Diffusion with Bilateral Learning" by Nikolaos Vettas

93-25 "Short-Run Independence of Monetary Policy under Pegged Exchange Rates and Effects of Money on Exchange Rates and Interest Rates" by Lee E. Ohanian and Alan C. Stockman

93-26 "Majority Rule and the Public Provision of Health Care" by Miguel Gouveia

93-27 "Pareto Improving Financial Innovation in Incomplete Markets" by David Cass and Alessandro Citanna

93-28 "On the Dynamic Selection of Mechanisms as Clubs for the Provision of a Public Project" by Roger D. Lagunoff

94-01 "Expected Utility and Case-Based Reasoning" by Akihiko Matsui

94-02 "Sequential Stratified Sampling" by Edward J. Green and Ruilin Zhou

94-03 "Bargaining, Boldness and Nash Outcomes" by Simon Grant and Atsushi Kajii

94-04 "Learning and Strategic Pricing" by Dirk Bergemann and Juuso Valimaki

94-05 "Evolution in Mechanisms for Public Projects" by Roger D. Lagunoff and Akihiko Matsui (previous version 93-14)

94-06 "Constrained Suboptimality in Incomplete Markets: A General Approach and Two Applications" by Alessandro Citanna, Atsushi Kajii and Antonio Villanacci

- 94-07 "Pareto Improving Financial Innovation in Incomplete Markets" by David Cass and Alex Citanna (previous version 93-27)
- 94-08 "Commodity Money Under Private Information" by Yiting Li
- 94-09 "Generic Local Uniqueness in the Walrasian Model: A Pedagogical Note" by Marcos de Barros Lisboa
- 94-10 "Bargaining-Induced Transaction Demand for Fiat Money" by Merwan Engineer and Shouyong Shi
- 94-11 "Politico-Economic Equilibrium and Economic Growth" by Per Krusell, Vincenzo Quadrini and José- Víctor Ríos-Rull
- 94-12R "On the Evolution of Pareto Optimal Behavior in Repeated Coordination Problems" by Roger D. Lagunoff
- 94-13 "Evolution and Endogenous Interactions" by George J. Mailath, Larry Samuelson and Avner Shaked
- 94-14R "How Proper is Sequential Equilibrium?" by George J. Mailath, Larry Samuelson and Jeroen M. Swinkels
- 94-15 "Common p-Belief: The General Case" by Atsushi Kajii and Stephen Morris
- 94-16 "Impact of Public Announcements on Trade in Financial Markets" by Stephen Morris and Hyun Song Shin
- 94-17 "Payoff Continuity in Incomplete Information Games and Almost Uniform Convergence of Beliefs" by Atsushi Kajii and Stephen Morris
- 94-18 "Public Goods and the Oates Decentralisation Theorem" by Julian Manning
- 94-19 "The Rationality and Efficacy of Decisions under Uncertainty and the Value of an Experiment" by Stephen Morris and Hyun Song Shin
- 94-20 "Does Rational Learning Lead to Nash Equilibrium in Finitely Repeated Games?" by Alvaro Sandroni
- 94-21 "On the Form of Transfers to Special Interests" by Stephen Coate and Stephen Morris
- Revised and final version is forthcoming in the Journal of Political Economy
- 94-22 "Specialization of Labor and the Distribution of Income" by Akihiko Matsui and Andrew Postlewaite
- 95-01 "Financial Innovation and Expectations" by Alessandro Citanna and Antonio Villanacci
- 95-02 "An Economic Model of Representative Democracy" by Tim Besley and Stephen Coate
- 95-03 "The Revelation of Information and Self-Fulfilling Beliefs" by Jayasri Dutta and Stephen Morris

- 95-04 “Justifying Rational Expectations” by Stephen Morris
- 95-05 “Co-operation and Timing” by Stephen Morris
- 95-06 “Statistical Discrimination, Affirmative Action, and Mismatch” by Jaewoo Ryoo
- 95-07 “Sufficiently Specialized Economies have Nonempty Cores” by Roger D. Lagunoff
- 95-08 “Necessary and Sufficient Conditions for Convergence to Nash Equilibrium: The Almost Absolute Continuity Hypothesis” by Alvaro Sandroni
- 95-09 “Budget-constrained Search” by Richard Manning and Julian Manning
- 95-10 “Efficient Policy Choice in a Representative Democracy: A Dynamic Analysis” by Timothy Besley and Stephen Coate
- 95-11 “The Sequential Regularity of Competitive Equilibria and Sunspots” by Atsushi Kajii
- 95-12 “Generic Existence of Sunspot Equilibria: The Case of real Assets” by Piero Gottardi and Atsushi Kajii
- 95-13 “Speculative Investor Behavior and Learning” by Stephen Morris
- 95-14 “Incorporating Concern for Relative Wealth into Economic Models” by Harold L. Cole, George J. Mailath and Andrew Postlewaite
- 95-15 “An ‘Anti-Folk Theorem’ for a Class of Asynchronously Repeated Games” by Roger Lagunoff and Akihiko Matsui